

**Obstetric Anesthesia During the COVID-19 Pandemic**

Melissa Bauer, DO<sup>1</sup>; Kyra Bernstein, MD<sup>2</sup>; Emily Dinges, MD<sup>3</sup>; Carlos Delgado, MD<sup>3</sup>; Nadir El-Sharawi, MD<sup>4</sup>; Pervez Sultan, MD<sup>5</sup>; Jill M. Mhyre, MD<sup>4</sup>; Ruth Landau, MD<sup>2</sup>

<sup>1</sup>Department of Anesthesiology, University of Michigan, Ann Arbor, MI

<sup>2</sup>Department of Anesthesiology, Columbia University College of Physicians and Surgeons, New York, NY

<sup>3</sup>Department of Anesthesiology & Pain Medicine, University of Washington, Seattle, WA

<sup>4</sup>Department of Anesthesiology, University of Arkansas for Medical Sciences, Little Rock, AR

<sup>5</sup>Department of Anesthesia, Stanford University, Stanford, CA

**Corresponding Author:** Ruth Landau, MD, Department of Anesthesiology, Columbia University College of Physicians and Surgeons, New York, NY

**Financial Disclosures:** None

**Conflicts of Interest:** None

**Word Count:** Abstract-110; Introduction-78; Discussion-NA; Overall-3128

**Abbreviated title:** Obstetric Anesthesia and COVID-19

**Author Contributions:**

Melissa Bauer: This author helped write the manuscript and edit for critical content.

Kyra Bernstein: This author helped write the manuscript and edit for critical content.

Emily Dinges: This author helped write the manuscript and edit for critical content.

Carlos Delgado: This author helped write the manuscript and edit for critical content.

Nadir El-Sharawi: This author helped write the manuscript and edit for critical content.

Pervez Sultan: This author helped write the manuscript and edit for critical content.

Jill Mhyre: This author helped write the manuscript and edit for critical content.

Ruth Landau: This author helped write the manuscript and edit for critical content.

ACCEPTED

**Abstract:**

With increasing numbers of Coronavirus Disease 2019 (COVID-19) cases due to efficient human-to-human transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in the United States, preparation for the unpredictable setting of labor and delivery is paramount. The priorities are two-fold in the management of obstetric patients with COVID-19 infection or persons under investigation (PUI): (1) caring for the range of asymptomatic to critically ill pregnant and postpartum women; (2) protecting health care workers and beyond from exposure during the delivery hospitalization (health care providers, personnel, family members). The goal of this review is to provide evidence-based recommendations, or expert opinion when evidence is limited, for anesthesiologists caring for pregnant women during the COVID-19 pandemic, with a focus on preparedness and best clinical obstetric anesthesia practice.

**Glossary of Terms:**

COVID-19: Coronavirus Disease 2019

SARS-CoV-2 (COVID-19): severe acute respiratory syndrome coronavirus 2

RT-PCR: real-time reverse transcriptase–polymerase chain reaction

PUI: person under investigation for COVID-19 infection

ARDS: Acute Respiratory Distress Syndrome (ARDS)

SARS: severe acute respiratory syndrome

PaCO<sub>2</sub>: partial pressure of carbon dioxide

NSAIDs: nonsteroidal anti-inflammatory drugs

## **Introduction**

The management of obstetric patients infected with Coronavirus Disease 2019 (COVID-19) due to human-to-human transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) requires quite unique considerations, from caring for critically ill pregnant and postpartum women to protecting health care workers from exposure during the delivery hospitalization (health care providers, personnel, family members and beyond). The goal of this review is to provide evidence-based recommendations, or expert opinion when evidence is limited, for anesthesiologists caring for pregnant women during the COVID-19 pandemic, with a focus on preparedness and best clinical obstetric anesthesia practice.

## **Clinical Manifestation of COVID-19 Infection in Pregnancy**

In principle, the clinical characteristics reported in pregnant women with confirmed COVID-19 infection in China have been consistent with those reported among non-pregnant adults, with better maternal and neonatal outcomes with COVID-19 infection compared with the 2002-2003 severe acute respiratory syndrome (SARS) outbreak from SARS-CoV-1 infection.<sup>1-3</sup>

The signs and symptoms of COVID-19 infection in a large data set in non-pregnant patients from China were fever (99%), fatigue (70%), cough (59%), shortness of breath (31%), myalgias (35%), headache (6.5%), sore throat (17%), diarrhea (10%), nausea (10%), and vomiting (4%).<sup>4</sup> An additional manifestation noted among patients with COVID-19 infection is the sudden loss (or reduction) of the sense of smell and taste, which is currently recommended by the American Academy of Otolaryngology - Head and Neck Surgery as part of screening for COVID-19 infection.<sup>5</sup>

In pregnancy, presentation of COVID-19 infection appears similar, but many of these non-specific symptoms may be attributed to symptoms of pregnancy and labor.<sup>2</sup> For example, signs of latent labor may include myalgias and diarrhea; preeclampsia can present with

headache; shortness of breath is perceived during pregnancy and labor; and chorioamnionitis may cause tachycardia and fever—leading clinicians to overlook COVID-19 infection as a possible diagnosis. In addition, women infected with COVID-19 may be asymptomatic until their admission in labor and beyond,<sup>6</sup> which in itself poses a significant risk of exposure for their family members (including the newborn) and all providers involved in their clinical care.

### **Considerations for Testing on Admission**

Screening criteria for COVID-19 infection usually includes the following: (1) fever, (2) cough or shortness of breath, (3) diarrhea, and (4) any possible exposure to COVID-19. However, because women with COVID-19 infection may be asymptomatic at the time of admission, and because some may present with overlapping pregnancy symptoms, universal screening may miss pregnant women infected with SARS CoV2 in communities with a high prevalence or high projected infection rate (e.g., New York, New Orleans, Detroit, Chicago, Miami).<sup>7,8</sup> Universal testing with real-time reverse transcriptase–polymerase chain reaction (RT-PCR) tests for SARS-CoV-2 viral RNA may improve case detection in high prevalence communities. However, current assays may return false negative results if the viral load is low or if specimen collection is incomplete.

The goals of COVID-19 testing specific to pregnant patients admitted to labor and delivery units are two-fold: (1) to prevent vertical transmission and ensure separation of the neonate after birth, and (2) to protect health care workers by ensuring use of appropriate personal protective equipment (PPE). Besides the unclear sensitivity of RT-PCR testing, the time for nucleic acid detection varies between 6 to 8 hours or longer depending on availability. Therefore, management of women on Labor and Delivery Units located in a community with a high prevalence of COVID-19 infection should err on the side of caution.

For purposes of clinical management and PPE use, women may therefore be categorized as follows (1) COVID-19 negative, (2) asymptomatic, (3) symptomatic (PUI), and (4) personally positive for COVID-19 testing. This information should be made available to all health care providers and updated at all times, as it may change during the course of labor (from asymptomatic to symptomatic, or if tested, once the result becomes available).

Women who are COVID-19 positive (or high risk PUI) should ideally be placed in an isolation room. Airborne infection isolation rooms (single-patient negative-pressure rooms with a minimum of 6 air changes per hour), if available, should be used if performance of aerosolizing procedures is anticipated. In general, isolation rooms suitable for droplet and contact precautions are recommended.<sup>9</sup> Strategies for exposure mitigation and cohorting, as well as considerations for transportation of patients who are persons under investigation (PUI) or COVID-19 should follow the same recommendations as for general patient cases.<sup>10</sup>

### **Systems Preparation**

A multidisciplinary team of anesthesiologists, obstetricians, labor and delivery nurses, neonatologists, critical care experts, infectious disease and infection control experts, employee health services, environmental health services, and telemedicine services should create and implement protocols to support the management of patients with COVID-19 infection in the setting of a Labor and Delivery Unit. A side-by-side comparison of recommendations from many professional societies for labor and delivery units is presented in Table 1. For institutions with multiple labor and delivery sites, consideration should be given to designating one institution to care for patients with COVID-19 infection. This proved useful in managing patients during the SARS epidemic and for cases in the recent COVID-19 outbreak in Wuhan, China.<sup>11-13</sup>

Resource allocation within the Labor and Delivery Unit as well as other units (including Intensive Care Unit) should be proactively addressed. It is imperative to establish a back-up team to care for patients without COVID-19 infection due to the time-intensive tasks of donning/doffing PPE, transporting the patient, providing anesthetic care, and performing surgery in patients with active COVID-19 infection.

From a logistical standpoint, a designated operating room within the Labor and Delivery Unit should be prepared. Dedicated trays (or carts) containing the most commonly used supplies and drugs for both neuraxial labor analgesia and cesarean delivery should be available, to minimize traffic and contamination of anesthesia workstations and other anesthesia equipment.

### **Patient Evaluation and Monitoring**

A pregnant woman who is a PUI or COVID-19 positive should be evaluated (limiting unnecessary encounters), including vital signs, physical examination, and review of laboratory tests (complete blood count, comprehensive metabolic panel, and arterial blood gas, if needed) to assess appropriate level of care and monitoring plan for potential deterioration. Early multidisciplinary collaboration should be arranged to determine level of care, fetal monitoring, and delivery plan. Discussion of the risks and benefits for administering steroids for fetal lung maturity, magnesium for neuroprotection, and indomethacin for tocolysis should be addressed, since there is concern those drugs may worsen COVID-19 infection (Table 2).<sup>9</sup>

Avoiding urgent cesarean delivery is essential, to reduce the risk for general anesthesia and provider exposure during uncontrolled transfers to the operating room. Therefore, ongoing assessment of both maternal and fetal status are key to balance risks of prolonged labor versus cesarean delivery. It is unclear whether uterine decompression improves maternal respiratory status and how the potential benefit balances against the known operative risks in the setting of

COVID-19. On the other hand, prolonged maternal hypoxemia may ultimately cause fetal acidemia, leading to a more urgent cesarean delivery.<sup>9</sup>

Routine monitoring should include frequent vital signs (tailored to the current clinical status and adjusted as necessary) with the addition of continuous pulse oximetry and strict input and output measurements to assure fluid restriction. Pulse oximetry goal should be an oxygen saturation  $\geq 95\%$ . Early warning criteria systems specific for obstetric patients may aid in early detection and prompt escalation of care.<sup>14</sup>

Women requiring supplemental oxygen, who develop increasing oxygen requirements or worsening hypoxia ( $SpO_2 < 95\%$ ), should have prompt arterial blood gas analysis with frequent clinical reassessment to guide the requirement for escalation of care and mechanical ventilation. High flow nasal oxygen or non-invasive ventilation may be considered as temporizing measures but are generally discouraged due to the potential for greater aerosolization. In addition, increasing oxygen requirements serve as a marker of disease progression, with increasing risk of atelectasis and pulmonary consolidation. It is recommended to perform early endotracheal intubation in a controlled manner minimizing exposure to health care workers and equipment with airborne precautions in an urgent/emergent situation.<sup>15</sup>

One of the frequent complications of patients with COVID-19 is Acute Respiratory Distress Syndrome (ARDS). Ventilator management strategies for ARDS involve lung-protective strategies, such as low tidal volumes (6 mL/kg using predicted body weight), plateau pressure  $< 30$  cm H<sub>2</sub>O, and the combined use of reduced FiO<sub>2</sub> with increases in positive end expiratory pressure (PEEP) to maintain a PaO<sub>2</sub> of 65-90 mm Hg.<sup>16</sup> Useful ventilator titration techniques using the ARDSnet ventilator protocol can be found here:

[http://www.ardsnet.org/files/ventilator\\_protocol\\_2008-07.pdf](http://www.ardsnet.org/files/ventilator_protocol_2008-07.pdf).<sup>17</sup> Pregnant patients have a physiological decrease in PaCO<sub>2</sub>, and it is recommended to maintain a PaCO<sub>2</sub> of 28-32 mm Hg



with ventilation to augment off-loading of oxygen to the fetus. However, the priority is maintaining oxygenation with low tidal volumes and PEEP, and this strategy may not allow for maintaining the physiologic PaCO<sub>2</sub> in pregnancy. Multidisciplinary discussion should determine the fetal monitoring and delivery plan during mechanical ventilation.

### **Laboring Patients with Active COVID-19 Infection**

Neuraxial labor analgesia remains a mainstay of obstetric care even with concurrent COVID-19 infection. In fact, early epidural placement is desirable to avoid exacerbation of respiratory symptoms with labor pain, and to reduce the likelihood of general anesthesia if intrapartum cesarean delivery becomes needed.

The benefits of neuraxial analgesia in the setting of COVID-19 pneumonia are two-fold: (1) for the patient it will help avoid any exacerbation of respiratory status with intubation and mechanical ventilation, (2) for health care providers, it reduces the risks associated with aerosol exposure and transmission of COVID-19 infection during intubation and extubation if general anesthesia is provided.

The risk of COVID-19 exposure for the anesthesiologist during neuraxial labor analgesia placement is presumably low, since this is not an aerosol-generating procedure. All health care workers in the room should wear contact (impervious gown and gloves) and droplet (surgical mask and eye protection) precautions. The patient should wear a surgical mask at all times to limit droplet spread, and the number of personnel present during placement of neuraxial labor analgesia should be minimized, but with assistance readily available. Several strategies may minimize contamination of equipment and COVID-19 exposure in anesthesiologists, while also minimizing the consumption of PPE (Box 1 and Figure 1).

A parturient who is symptomatic (PUI), or COVID-19 positive, should have a complete blood count prior to neuraxial analgesia placement. Early studies from China suggested that thrombocytopenia may be associated with COVID-19 infection; in a cohort of 1099 patients 36.2% had thrombocytopenia ( $< 150,000 \times 10^6/L$ ).<sup>18</sup> A meta-analysis of 1779 patient with COVID-19 infection observed that platelet counts are lower in patients with more severe disease.<sup>19</sup> Though less common, a platelet count less than  $100,000 \times 10^6/L$  can occur; three studies of 243 patients reported a total of 6.6% patients with that level of thrombocytopenia.<sup>20-22</sup>

We suggest a platelet count on admission, without the need to check serial counts prior to needle placement unless there is a major change in clinical symptoms. It is generally safe to perform neuraxial procedures at platelet counts of  $70,000 \times 10^6/L$  or above,<sup>23</sup> and given the rare risk of spinal/epidural hematoma and the much higher risk of respiratory compromise with general anesthesia, neuraxial procedures at even lower platelet counts should be considered.

While theoretically possible, the risk of epidural or subarachnoid space seeding with viremic blood, causing encephalitis or meningitis is exceedingly rare. At the time of this writing, there are 77 pregnant women reported in the literature who received uneventful neuraxial procedures for cesarean or vaginal delivery (4 combined spinal-epidural, 27 epidural, 46 spinal procedures; Table 3).<sup>6,24-28</sup> Spinal, combined spinal-epidural, epidural procedures all are acceptable, and no technique confers more risk than the other based on the literature available. None of the patients experienced neurologic sequelae.

Current recommendations on the use of nitrous oxide (Entonox) for labor analgesia suggest “there is insufficient information about the cleaning, filtering, and potential aerosolization of nitrous oxide in the setting of COVID-19.” Individual Labor and Delivery Units should consider suspending use.<sup>9</sup> Additionally, the practice of high flow oxygen for fetal distress does not improve fetal outcomes and should be suspended due to the risk of aerosolization.<sup>9</sup>

## **Anesthesia for Cesarean Delivery**

In reports from China, most women with a diagnosis of COVID-19 infection underwent a cesarean delivery.<sup>24</sup> In the absence of universal testing and rapid availability of results, COVID-19 status may not necessarily be known at the time of cesarean delivery. The baseline failure rate for conversion of labor epidural analgesia to cesarean delivery anesthesia is 5%.<sup>29</sup> Urgent intrapartum cesarean delivery represents an important risk factor for failed conversion from intrapartum neuraxial labor analgesia to cesarean delivery anesthesia—therefore ongoing communication with the obstetricians is crucial to allow safe transfer to the operating room, and adequate time for initiation of surgical block to avoid general anesthesia.<sup>29</sup> To minimize the risk of exposure during urgent endotracheal intubation, airborne protection (N95 respirator mask) is recommended for all providers in the operating room unless the patient is known to be COVID-19 negative.

A publication from Wuhan, China describing outcomes in 17 COVID-19 positive women undergoing cesarean delivery concluded that “excessive hypotension” occurred in 12 of 14 cases with epidural anesthesia in comparison with the 3 women who had received general anesthesia; however, information about the blood pressure trends and description of the use of vasopressors is not reported.<sup>25</sup> A larger case series of 49 patients receiving spinal anesthesia (45 for cesarean delivery and 4 for orthopedic procedures) was well-tolerated with stable blood pressure.<sup>28</sup>

In our early experience, maternal hypotension during cesarean delivery with epidural or spinal anesthesia has not been noted, most likely because prevention of hypotension with phenylephrine is part of our routine clinical practice. Along with anti-hypotensive medication, anti-emetic medication should also be administered. However, we recommend using an alternative to dexamethasone in a PUI or patient with known COVID-19 infection given the risk

of worsening clinical severity.<sup>30</sup> Specific considerations for medication use in PUI or COVID-19 positive patients during labor, delivery and the postpartum period are described in Table 2.

### **Vertical Transmission**

Current understanding is that there is little evidence for vertical transmission in women who develop COVID-19 pneumonia in late pregnancy.<sup>2,31-35</sup> However, cases of possible *in utero* infection seem to be emerging, including a recent report of a neonate born to a COVID-19 infected mother. This suggests *in utero* infection during the 23 days between maternal infection and delivery 23 days later, supported by elevated IgM antibodies, which are not transferred to the fetus via the placenta.<sup>36-39</sup> Serological testing of virus specific IgG and IgM antibodies may alternatively be used if RT-PCR testing is not available or if RT-PCR seems to be yielding a false-negative result.<sup>40</sup>

### **Postpartum Considerations**

Postpartum considerations for parturients with COVID-19 infection include adequate management of usual postpartum issues (postpartum hemorrhage, pain, hemodynamic status), as well as judicious fluid management, surveillance for respiratory decompensation, and early involvement of subspecialty care as needed. Appropriate isolation of mother and child on the postpartum unit is also recommended.<sup>41</sup>

#### *Postpartum hemorrhage*

In the setting of postpartum hemorrhage due to uterine atony, carboprost tromethamine (Hemabate®) followed by endotracheal intubation was reported to have precipitated immediate and prolonged bronchospasm in a patient who was subsequently found to be COVID-19 positive.<sup>6,42</sup> Oxytocin and methylergonovine as a second-line choice may be preferred, due to the potential for bronchospasm with carboprost tromethamine (Hemabate®), and aerosolization of viral particles during bronchospasm management.

### *Post-partum pain management*

It has been posited that nonsteroidal anti-inflammatory drugs (NSAIDs) for management of symptoms suggestive of COVID-19 infection may worsen the clinical course of COVID-19 patients; however, this remains controversial and robust data are lacking. At this point, for women who are asymptomatic or mildly symptomatic with pain not well-controlled with acetaminophen, NSAIDs can continue to be used, as the alternative of opioids likely poses more clinical risks.<sup>9</sup>

### *Management of post-dural puncture headache*

There are no reported cases of accidental dural puncture resulting in post-dural puncture headache (PDPH) in a patient with a COVID-19 infection, and consequently, no available guidance. Similar to usual care, conservative measures should be initially provided.

Usual contraindications to the performance of an epidural blood patch (e.g., fever, thrombocytopenia, or other coagulation issues) should apply in a COVID-19 patient. Mitigating the risk of a serious neurologic complications with untreated PDPH<sup>43</sup> versus that of viral seeding in the epidural space with an epidural blood patch will require a case-by-case approach. Postponing the epidural blood patch is recommended in women who are actively ill. Individual assessment of the benefits and risk should be assessed and shared decision-making should be engaged with the patient prior to proceeding.

Because a nasal sphenopalatine ganglion (SPG) block is likely an aerosol generating procedure due to the injection/insertion directly into the nasal cavity, it should be avoided to minimize the risk of transmission to health care workers.

## Conclusions

Key points emerging in the past weeks from the literature and our experience in Labor and Delivery Units in the United States are that pregnant women may be asymptomatic upon admission in labor, and that symptoms of COVID-19 infection may initially be missed or obscured if chorioamnionitis is suspected during labor. Although most women with COVID-19 infection will not present with pneumonia and respiratory decompensation during labor, escalation of care and advanced critical care resources may become necessary in the postpartum period.

In fact, most of the considerations surrounding management of the parturient with suspicion of or known COVID-19 infection include not only best strategies to ensure safe care for the parturient but also to prevent health care worker exposure to SARS-CoV-2 and contracting COVID-19.

Anesthesiologists are deemed at significant risk of viral exposure during endotracheal intubations of COVID-19 infected patients, and all strategies should be applied to avoid general anesthesia in women who are either untested or known to be COVID-19 positive.

Early neuraxial labor analgesia is strongly recommended to ensure availability of neuraxial anesthesia in the event of an intrapartum cesarean delivery, and spinal anesthesia should be provided if needed. If deemed necessary and not avoidable, provision of general anesthesia should follow general recommendations for intubation and extubation in the setting of COVID-19 infected patients.

The changes in workflow that result from the need to ensure adequate PPE (contact/droplet protection for non-aerosolizing procedures such as epidural placement, or airborne protection for cesarean deliveries due to the possible conversion to general anesthesia) are considerable and require thorough planning and preparedness.

Close communication around COVID-19 status of all patients admitted to the Labor and Delivery Unit is essential, and anticipation of emergencies is of the essence. Overall, providing best clinical care for pregnant and postpartum women with COVID-19 infection also must take into account strategies to prevent health care worker exposure to SARS-CoV-2 and contracting COVID-19.

ACCEPTED

## References

1. Qiao J. What are the risks of COVID-19 infection in pregnant women? *Lancet* 2020;395:760-2.
2. Chen H, Guo J, Wang C et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* 2020;395:809-15.
3. Liu Y, Chen H, Tang K, Guo Y. Clinical manifestations and outcome of SARS-CoV-2 infection during pregnancy. *J Infect* 2020.
4. Wang D, Hu B, Hu C et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA* 2020.
5. American Academy of Otolaryngology - Head and Neck Surgery. COVID-19 Anosmia Reporting Tool for Clinicians. 2020. <https://www.entnet.org/content/reporting-tool-patients-anosmia-related-covid-19>. Accessed March 28, 2020.
6. Breslin N, Baptiste C, Miller R et al. COVID-19 in pregnancy: early lessons. *Am J Obstet Gynecol MFM* 2020.
7. Coronavirus Disease 2019 (Covid-19): Cases in the U.S. Centers for Disease Control and Prevention. 2020. <https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/cases-in-us.html>. Accessed March 27, 2020.
8. Institute for Health Metrics and Evaluation (IHME). COVID-19 Projections. 2020. <http://www.healthdata.org/data-visualization/covid-19-us-state-state-projections>. Accessed March 27, 2020.



9. Society for Maternal-Fetal Medicine, Society for Obstetric and Anesthesia and Perinatology. Labor and Delivery COVID-19 Considerations. 2020.  
[https://s3.amazonaws.com/cdn.smfm.org/media/2277/SMFM-SOAP\\_COVID\\_LD\\_Considerations\\_3-27-20\\_\(final\)\\_PDF.pdf](https://s3.amazonaws.com/cdn.smfm.org/media/2277/SMFM-SOAP_COVID_LD_Considerations_3-27-20_(final)_PDF.pdf). Accessed March 27, 2020.
10. Center for Disease Control and Prevention. Interim Infection Prevention and Control Recommendations for Patients with Suspected or Confirmed Coronavirus Disease 2019 (COVID-19) in Healthcare Settings. 2020. [https://www.cdc.gov/coronavirus/2019-ncov/infection-control/control-recommendations.html#Patient\\_Placement](https://www.cdc.gov/coronavirus/2019-ncov/infection-control/control-recommendations.html#Patient_Placement). Accessed March 27, 2020.
11. Zhang H-F, Bo L-L, Lin Y et al. Response of Chinese Anesthesiologists to the COVID-19 Outbreak. *Anesthesiology* 2020.
12. Haines CJ, Chu YW, Chung TK. The effect of Severe Acute Respiratory Syndrome on a hospital obstetrics and gynaecology service. *BJOG : an international journal of obstetrics and gynaecology* 2003;110:643-5.
13. Maxwell C, McGeer A, Tai KFY et al. Management Guidelines for Obstetric Patients and Neonates Born to Mothers With Suspected or Probable Severe Acute Respiratory Syndrome (SARS). *J Obstet Gynaecol Can* 2009;31:358-64.
14. Mhyre JM, D'Oria R, Hameed AB et al. The maternal early warning criteria: a proposal from the national partnership for maternal safety. *J Obstet Gynecol Neonatal Nurs* 2014;43:771-9.
15. Jin Y-H, Cai L, Cheng Z-S et al. A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus (2019-nCoV) infected pneumonia (standard version). *Mil Med Res* 2020;7:4.

16. Duarte AG. ARDS in pregnancy. *Clin Obstet Gynecol* 2014;57:862-70.
17. NIH-NHLBI ARDS Network. NIH NHLBI ARDS Clinical Network Mechanical Ventilation Protocol Summary. 2008.  
[http://www.ardsnet.org/files/ventilator\\_protocol\\_2008-07.pdf](http://www.ardsnet.org/files/ventilator_protocol_2008-07.pdf). Accessed March 28.
18. Guan W-j, Ni Z-y, Hu Y et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine* 2020.
19. Lippi G, Plebani M, Henry BM. Thrombocytopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: A meta-analysis. *Clinica Chimica Acta* 2020;506:145-8.
20. Huang C, Wang Y, Li X et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 2020;395:497-506.
21. Liu Y, Yang Y, Zhang C et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Science China Life Sciences* 2020;63:364-74.
22. Zhou F, Yu T, Du R et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* 2020;395:1054-62.
23. Lee LO, Bateman BT, Kheterpal S et al. Risk of Epidural Hematoma after Neuraxial Techniques in Thrombocytopenic Parturients: A Report from the Multicenter Perioperative Outcomes Group. *Anesthesiology* 2017;126:1053-63.
24. Bauer ME, Chiware R, Pancaro C. Neuraxial procedures in COVID-19 positive parturients: a review of current reports. *Anesth Analg* 2020.

25. Chen R, Zhang Y, Huang L et al. Safety and efficacy of different anesthetic regimens for parturients with COVID-19 undergoing Cesarean delivery: a case series of 17 patients. *Can J Anaesth* 2020;1-9.
26. Song L, Xiao W, Ling K et al. Anesthetic Management for Emergent Cesarean Delivery in a Parturient with Recent Diagnosis of Coronavirus Disease 2019 (COVID-19): A Case Report. *Transl Perioper Pain Med* 2020;7:234-7.
27. Xia H, Zhao S, Wu Z et al. Emergency Caesarean delivery in a patient with confirmed coronavirus disease 2019 under spinal anaesthesia. *Br J Anaesth* 2020.
28. Zhong Q, Liu YY, Luo Q et al. Spinal anaesthesia for patients with coronavirus disease 2019 and possible transmission rates in anaesthetists: retrospective, single-centre, observational cohort study. *British Journal of Anaesthesia* 2020.
29. Bauer M, Kountanis J, Tsen L et al. Risk factors for failed conversion of labor epidural analgesia to cesarean delivery anesthesia: a systematic review and meta-analysis of observational trials. *Int J Obstet Anesth* 2012;21:294-309.
30. Society for Obstetric and Anesthesia and Perinatology. Interim Considerations for Obstetric Anesthesia Care related to COVID19. 2020. [https://soap.org/wp-content/uploads/2020/03/SOAP\\_COVID-19\\_Obstetric\\_Anesthesia\\_Care\\_032320.pdf](https://soap.org/wp-content/uploads/2020/03/SOAP_COVID-19_Obstetric_Anesthesia_Care_032320.pdf). Accessed March 27, 2020.
31. Fan C, Lei D, Fang C et al. Perinatal Transmission of COVID-19 Associated SARS-CoV-2: Should We Worry? *Clin Infect Dis* 2020.
32. Li Y, Zhao R, Zheng S et al. Lack of Vertical Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, China. *Emerg Infect Dis* 2020;26.

33. Schwartz DA. An Analysis of 38 Pregnant Women with COVID-19, Their Newborn Infants, and Maternal-Fetal Transmission of SARS-CoV-2: Maternal Coronavirus Infections and Pregnancy Outcomes. *Arch Pathol Lab Med* 2020.
34. Zhu H, Wang L, Fang C et al. Clinical analysis of 10 neonates born to mothers with 2019-nCoV pneumonia. *Transl Pediatr* 2020;9:51-60.
35. Stower H. Lack of maternal–fetal SARS-CoV-2 transmission. *Nat Med* 2020;26:312.
36. Dong L, Tian J, He S et al. Possible Vertical Transmission of SARS-CoV-2 From an Infected Mother to Her Newborn. *JAMA* 2020.
37. Zeng H, Xu C, Fan J et al. Antibodies in Infants Born to Mothers With COVID-19 Pneumonia. *JAMA* 2020.
38. Kimberlin DW, Stagno S. Can SARS-CoV-2 Infection Be Acquired In Utero?: More Definitive Evidence Is Needed. *JAMA* 2020.
39. Wang S, Guo L, Chen L et al. A Case Report of Neonatal 2019 Coronavirus Disease in China. *Clin Infect Dis* 2020.
40. Dong X, Cao Y-y, Lu X-x et al. Eleven Faces of Coronavirus Disease 2019. *Allergy* 2020.
41. Rasmussen SA, Smulian JC, Lednicky JA et al. Coronavirus Disease 2019 (COVID-19) and Pregnancy: What obstetricians need to know. *Am J Obstet Gynecol* 2020.
42. Landau R, Bernstein K, Mhyre J. Lessons learned from first COVID-19 cases in the United States. *Anesth Analg* 2020;In Press.
43. Guglielminotti J, Landau R, Li G. Major Neurologic Complications Associated With Postdural Puncture Headache in Obstetrics: A Retrospective Cohort Study. *Anesth Analg* 2019;129:1328-36.

44. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *The Lancet Respiratory Medicine* 2020.

ACCEPTED

**Figure Legends.**

**Figure 1. Set-up for Operating Room and Labor Epidural Analgesia**

ACCEPTED

**Table 1. Professional Society Recommendations for Labor and Delivery**

<b>Comparison of Recommendations for Intra-Partum Anesthetic Care of Known or Suspected COVID-19 Patients</b>							
	<b>SOAP</b>	<b>ACOG</b>	<b>SMFM-SOAP</b>	<b>SOGC</b>	<b>RANZCOG</b>	<b>RCOA-OAA</b>	<b>RCOG</b>
<b>Initial publication</b>	March 15, 2020	March 23, 2020	March 25, 2020	March 17, 2020	March 23, 2020	March 16, 2020	March 15, 2020
<b>Date updated</b>	March 23, 2020	March 26, 2020	March 27, 2020	March 26, 2020	March 29, 2020	March 27, 2020	March 28, 2020
<b>Isolation room</b>	Yes, preferably negative pressure	--	Yes, negative pressure for AGP	--	Yes	--	Yes
<b>Mask on patient</b>	--	Yes, surgical	Yes, surgical	Yes, surgical	Yes, surgical	Yes, surgical	Yes, surgical
<b>Visitors and support people</b>	Limit visitors and support people	Limit visitors and support people	Limit or eliminate visitors, special considerations apply	Yes, single asymptomatic	Yes, limited, asymptomatic	--	Yes, single asymptomatic, limit visitors to wards
<b>PPE for non-AGP, 1<sup>st</sup> stage labor</b>	Droplet and contact	Droplet and contact, airborne preferred	Droplet and contact	Droplet and contact	--	--	Droplet and contact
<b>PPE for vaginal delivery, 2<sup>nd</sup> stage labor</b>	Droplet and contact	Droplet and contact, airborne preferred	Airborne reasonable, consider surgical drapes as additional barrier	Droplet and contact, consider airborne if risk of sudden deterioration	--	--	Droplet and contact
<b>PPE neuraxial analgesia</b>	Droplet and contact	Droplet and contact, airborne preferred	--	--	--	Droplet and contact	--
<b>Neuraxial analgesia and anesthesia COVID-specific concerns</b>	Labor analgesia encouraged early; Performed by most experienced provider	--	Labor analgesia encouraged early	Labor analgesia encouraged early; Performed by most experienced provider	--	Labor analgesia encouraged early, suggestion to check platelet count	Labor analgesia encouraged early

<b>Use of nitrous oxide</b>	Consider suspending use	--	Consider suspending use	--	Yes, no evidence of AGP, use filter	Yes, no evidence of AGP, use filter	Yes, no evidence of AGP, use filter
<b>Use of post-operative NSAIDs</b>	Data lacking, likely safe if asymptomatic	Data lacking	Data lacking, continue to use in asymptomatic and mildly symptomatic patients	--	--	--	--
<b>Mode of delivery</b>	--	Per obstetric indications	--	Per obstetric indications	Per obstetric indications	--	Per obstetric indications, no water births
<b>Operative delivery</b>	--	Follow usual clinical indications	Not precluded, consider performing in OR	--	--	--	Individualized decision
<b>Elective cesarean or induction of labor with current COVID-19 infection</b>	--	--	--	--	Preferable to delay until completion of isolation period, if possible	--	Evaluate safety of delaying procedure
<b>PPE for cesarean delivery</b>	Anesthesia team: droplet, contact, ideally airborne; If GA planned: all staff wear airborne, minimize staff in room at time of intubation and extubation	Ideally airborne for all staff	Ideally airborne for all staff for all cases	Airborne advised; If GA planned: all staff wear airborne, minimize staff in room at time of intubation and extubation	--	Neuraxial planned (and low risk of GA): droplet and contact; Emergency cesarean: as above, risk assess need for airborne PPE; If GA planned: airborne PPE for all staff in room during intubation and extubation, minimize staff present, airborne for all staff if no time to allow for air turnover	Neuraxial planned (and low risk of GA): droplet, scrubbed staff waits outside of room until block successful; Emergency cesarean: droplet for neuraxial, airborne if risk of GA; If GA planned: airborne for all staff



**Table 2. Specific considerations for medication use in PUI or COVID-19 positive patients during labor, delivery and the postpartum period**

	<b>Considerations</b>	<b>Case context</b>	<b>Mechanism</b>
Oxygen <sup>9</sup>	The routine use of oxygen for fetal indications should be suspended	Overall, the use of oxygen for fetal indications is controversial	The use of high flow nasal cannula or face mask oxygen may be an aerosolizing procedure
Nitrous Oxide <sup>9</sup>	Discuss the relative risks and benefits of N <sub>2</sub> O for labor analgesia and consider suspending its use.	Overall, for all parturients	“There is currently insufficient information about the cleaning, filtering, and potential aerosolization of nitrous oxide in labor analgesia systems in the setting of COVID-19”
Remifentanyl/ fentanyl <sup>30</sup>	Consider avoiding the use of IV PCA opioids (remifentanyl/fentanyl) for labor analgesia	Women at risk for respiratory depression and opioid-induced nausea and vomiting	Opioid-induced respiratory depression increases the risk of sedation, respiratory depression and oxygen desaturation, and increases the risk for emergent airway instrumentation and aerosolizing procedures
Ketorolac/ ibuprofen <sup>9,44</sup>	For women who are asymptomatic or mildly symptomatic that require analgesic medication beyond acetaminophen, NSAIDs can continue to be used, as the alternative of opioids likely poses more clinical risks. For sick COVID-19 patients, consider avoiding NSAIDs	It has been suggested that the use of NSAIDs for management of COVID-19 symptoms may aggravate COVID-19 infection trajectory (although the evidence is not robust) The use of ACE inhibitors was suggested to increase the risk for COVID-19 infection	NSAIDs are associated with increased ACE2, to which COVID-19 binds

Dexamethasone <sup>9</sup>	Consider avoiding the use of dexamethasone for PONV prophylaxis	In all women undergoing cesarean delivery, alternative antiemetics should be administered to prevent vomiting	Prolonged exposure to high-dose steroids has been associated with worsening COVID-19 outcomes in the general population
Carboprost (Hemabate®) <sup>30</sup>	Consider avoiding the use of carboprost for treatment of uterine atony	For women at risk of bronchospasm, use alternative 2 <sup>nd</sup> line uterotonics	Prostaglandin F <sub>2</sub> alpha causes bronchoconstriction and pulmonary vasoconstriction
Magnesium sulfate <sup>9</sup>	Consider avoiding or as an alternative to usual dosing, a 4g bolus dose may be preferred in the setting of mild respiratory distress	For women with increasing oxygen requirement, the risk: benefit ratio should be considered before using magnesium for fetal neuroprotection, or for preeclampsia without severe features.	Magnesium sulfate has central nervous system and respiratory depressant effects

**Table 3.** Neuraxial procedures performed in pregnant women with COVID-19 infection

Author	n	Neuraxial procedure	Respiratory status at time of delivery	Febrile prior to procedure?	Neurologic sequelae
Bauer <sup>24</sup>	14	1 CSE, <sup>a</sup> 13 epidural	CT <sup>b</sup> or radiological evidence of pneumonia 14 (100%), 1 patient on maximal noninvasive ventilation at time of CD <sup>c</sup>	10 (71%)	None
Chen <sup>25</sup>	14	14 epidural	14 (100%) CT evidence of pneumonia	4 (29%)	None
Xia <sup>27</sup>	1	1 spinal	SpO <sub>2</sub> =92% on room air, CT evidence of pneumonia	1 (100%)	None
Song <sup>26</sup>	1	1 CSE	SpO <sub>2</sub> =86-90% on room air, CT evidence of pneumonia	1 (100%)	None
Breslin <sup>6e</sup>	2	2 CSE	Asymptomatic at the time of placement, found to have COVID-19 infection postpartum	0	None
Zhong <sup>28</sup>	45	Spinal	Mild symptoms, no cases of severe pneumonia	While hospitalized, 5 (11%) of entire cohort including 4 orthopedic patients	None

<sup>a</sup>CSE = combined spinal epidural, <sup>b</sup>CT = computed tomography, <sup>c</sup>CD = cesarean delivery, <sup>d</sup>SpO<sub>2</sub> = pulse oximetry, <sup>e</sup> Personal communication with Ruth Landau on March 29, 2020

**Figure 1**

**Signs to indicate precaution level**

**In OR: Airborne protection**  
(N-95 & fluid shield mask)



**In Labor Room: Contact/Droplet protection**



**Equipment readiness & access to supplies**

**OR preparedness**

Cover Anesthesia workstation



Buckets with supplies  
(uncovered here for display)



**Anesthesia machine**



Propofol  
Succinylcholine  
Rocuronium  
Lidocaine-Epi 2%



Ephedrine, phenylephrine  
Atropine, epinephrine, CaCl<sub>2</sub>  
Tranexamic acid  
Ondansetron, ketorolac



Spinal kit  
Chloragrep  
IV catheters  
IV tubing



Endotracheal tube  
Suction



HEPA filter  
Mask

ACCEPTED

**Box 1. Empirical strategies that may be implemented to minimize contamination of equipment and COVID-19 exposure of anesthesiologists to reduce the use of PPE**

1. Limit in-person encounters for pre-anesthesia evaluations by utilization of video - consultations (including for antenatal consultations for high-risk patients)
2. Limit encounters upon admission to the labor floor – consider using mounted iPads in each labor room for hourly evaluations
3. Limit the use of electronic devices or pens for written consents by use of electronically documented witnessed verbal consents
4. Avoid bringing into the patients’ room the epidural cart or tray - the required equipment (epidural kit) and drugs should be prepared and brought into the room in a bag before the procedure
5. Have the most experienced anesthesiologist perform the procedure to ensure adequate placement, and reduce the risk of accidental dural puncture that may require an epidural blood patch
6. Increase the dosing of neuraxial medications for labor analgesia (e.g. increasing the bupivacaine concentration from 0.0625% to 0.1%) or changing the setting of the programmed epidural intermittent bolus (e.g. increasing the volume from 5ml to 8ml, or decreasing the interval from every 45 minutes to 30 minutes) or adding neuraxial adjuvants (e.g. epidural clonidine) to minimize intrapartum breakthrough pain requiring epidural top-up
7. Round on parturients with video or phone calls into the patient’s room for hourly assessments of general status and effects of neuraxial analgesia
8. Ensure appropriate cleaning of all equipment in the room including the epidural pump and the on demand-button.
9. Limit encounters for postpartum rounds by utilization of video - consultations
10. Consider suspending prolonged patient-controlled epidural analgesia services (if applicable)